



SHARYN S.V.

APPLICATION OF THE FUNCTIONAL CALCULUS TO SOLVING OF INFINITE DIMENSIONAL HEAT EQUATION

In this paper we study infinite dimensional heat equation associated with the Gross Laplacian. Using the functional calculus method, we obtain the solution of appropriate Cauchy problem in the space of polynomial ultradifferentiable functions. The semigroup approach is considered as well.

Key words and phrases: infinite dimensional heat equation, Gross Laplacian, space of polynomial ultradifferentiable functions, space of polynomial ultradistributions.

Vasyl Stefanyk Precarpathian National University, 57 Shevchenka str., 76018, Ivano-Frankivsk, Ukraine
E-mail: sharyn.sergii@gmail.com

INTRODUCTION

The mathematical framework of white noise analysis, which was founded in works of Yu. Berezansky, Yu. Samoilenko [1] and T. Hida [5], is based on an infinite dimensional analogue of the Schwartz distribution theory.

In 1967 L. Gross [4] introduced Laplacian Δ_G on an abstract Wiener space as a natural infinite dimensional analogue of the finite dimensional Laplacian and studied potential theory associated with Δ_G . Within the white noise framework, the Gross Laplacian has been formulated by Kuo in [8] as a continuous linear operator acting on test white noise functions. The Gross Laplacian and appropriate Cauchy problem have been studied for example in [2, 9].

The aim of this work is to use the functional calculus constructed in [12] in order to solve the infinite dimensional heat equation associated with the Gross Laplacian.

1 PRELIMINARIES

1.1 Spaces of functions

Denote $\mathbb{Z}_+ := \{0\} \cup \mathbb{N}$ and $\partial^k := \partial^k / \partial t^k$. Fix any real $\beta > 1$. An infinitely differentiable function φ is called an ultradifferentiable function of the Gevrey class (see [7]) if for each segment $[\mu, \nu] \subset \mathbb{R}$ there exist constants $h > 0$ and $C > 0$ such that the inequality $\sup_{t \in [\mu, \nu]} |\partial^k \varphi(t)| \leq Ch^k k^{k\beta}$ holds for all $k \in \mathbb{Z}_+$. For a fixed $h > 0$ let us consider the subspace

$$\mathcal{G}_\beta^h[\mu, \nu] := \left\{ \varphi \in C^\infty : \text{supp } \varphi \subset [\mu, \nu], \|\varphi\|_{\mathcal{G}_\beta^h[\mu, \nu]} := \sup_{k \in \mathbb{Z}_+} \sup_{t \in [\mu, \nu]} \frac{|\partial^k \varphi(t)|}{h^k k^{k\beta}} < \infty \right\}.$$

Each subspace $\mathcal{G}_\beta^h[\mu, \nu]$ is a Banach space (see [7]) and all maps $\mathcal{G}_\beta^h[\mu, \nu] \hookrightarrow \mathcal{G}_\beta^l[\mu, \nu]$ with $h < l$ are compact inclusions. Consider the space

$$\mathcal{G}_\beta := \bigcup_{\mu < \nu, h > 0} \mathcal{G}_\beta^h[\mu, \nu], \quad \mathcal{G}_\beta \simeq \lim_{\mu < \nu, h > 0} \text{ind } \mathcal{G}_\beta^h[\mu, \nu],$$

of Gevrey ultradifferentiable functions with compact supports and endow it with topology of inductive limit with respect to above mentioned compact inclusions. Let \mathcal{G}'_β be its dual space of Roumieu ultradistributions.

Let $h > 0$ be any positive real and $\mu, \nu \in \mathbb{R}$ be any reals such that $\mu < \nu$. In the space of entire functions of exponential type we consider the subspace $E_\beta^h[\mu, \nu]$ of functions with the finite norm

$$\|\psi\|_{E_\beta^h[\mu, \nu]} := \sup_{k \in \mathbb{Z}_+} \sup_{z \in \mathbb{C}} \frac{|z^k \psi(z) e^{-H_{[\mu, \nu]}(\eta)}|}{h^k k^{k\beta}}, \quad \text{where } H_{[\mu, \nu]}(\eta) := \sup_{t \in [\mu, \nu]} t\eta.$$

Each space $E_\beta^h[\mu, \nu]$ is a Banach one, and all maps $E_\beta^h[\mu, \nu] \hookrightarrow E_\beta^{h'}[\mu', \nu']$ with $[\mu, \nu] \subset [\mu', \nu']$, $h < h'$, are compact inclusions. Consider the space

$$E_\beta := \bigcup_{\mu < \nu, h > 0} E_\beta^h[\mu, \nu], \quad E_\beta \simeq \lim_{\mu < \nu, h > 0} \text{ind } E_\beta^h[\mu, \nu],$$

and endow it with the topology of inductive limit with respect to above mentioned compact inclusions.

Consider the Fourier-Laplace transformation

$$\widehat{\varphi}(z) := (F\varphi)(z) = \int_{\mathbb{R}} e^{-itz} \varphi(t) dt, \quad \varphi \in \mathcal{G}_\beta, z \in \mathbb{C}.$$

Let $F' : E'_\beta \longrightarrow \mathcal{G}'_\beta$ be the adjoint mapping. It is known [13], that $F(\mathcal{G}_\beta) = E_\beta$ and $F'(E'_\beta) = \mathcal{G}'_\beta$.

1.2 Polynomial ultradifferentiable functions and polynomial ultradistributions

For any locally convex space \mathcal{X} , let $\mathcal{X}^{\widehat{\otimes} n}$, $n \in \mathbb{N}$, be the symmetric n th tensor degree of \mathcal{X} , completed in the projective tensor topology. For any $x \in \mathcal{X}$ we denote $x^{\otimes n} := \underbrace{x \otimes \cdots \otimes x}_n \in \mathcal{X}^{\widehat{\otimes} n}$, $n \in \mathbb{N}$. Set $\mathcal{X}^{\widehat{\otimes} 0} := \mathbb{C}$, $x^{\otimes 0} := 1 \in \mathbb{C}$.

To define the locally convex space $\mathcal{P}({}^n\mathcal{G}'_\beta)$ of n -homogeneous polynomials on \mathcal{G}'_β we use the canonical topological linear isomorphism $\mathcal{P}({}^n\mathcal{G}'_\beta) \simeq (\mathcal{G}'_\beta)^{\widehat{\otimes} n}$, described in [3]. We equip $\mathcal{P}({}^n\mathcal{G}'_\beta)$ with the locally convex topology \mathfrak{b} of uniform convergence on bounded sets in \mathcal{G}'_β . Set $\mathcal{P}({}^0\mathcal{G}'_\beta) := \mathbb{C}$. The space $\mathcal{P}(\mathcal{G}'_\beta)$ of all continuous polynomials on \mathcal{G}'_β is defined to be the complex linear span of all $\mathcal{P}({}^n\mathcal{G}'_\beta)$, $n \in \mathbb{Z}_+$, endowed with the topology \mathfrak{b} . Let $\mathcal{P}'(\mathcal{G}'_\beta)$ mean the strong dual of $\mathcal{P}(\mathcal{G}'_\beta)$. Elements of the spaces $\mathcal{P}(\mathcal{G}'_\beta)$ and $\mathcal{P}'(\mathcal{G}'_\beta)$ we call the polynomial test ultradifferentiable functions and polynomial ultradistributions, respectively.

Denote

$$\Gamma(\mathcal{G}_\beta) := \bigoplus_{n \in \mathbb{Z}_+} \text{fin } \mathcal{G}_\beta^{\widehat{\otimes} n} \subset \bigoplus_{n \in \mathbb{Z}_+} \mathcal{G}_\beta^{\widehat{\otimes} n} \quad \text{and} \quad \Gamma(\mathcal{G}'_\beta) := \bigotimes_{n \in \mathbb{Z}_+} \mathcal{G}'_\beta^{\widehat{\otimes} n}.$$

Note, that we consider only the case when the elements of direct sum consist of finite but not fixed number of addends. It is well known [11, 4.4], that $\langle \Gamma(\mathcal{G}'_\beta), \Gamma(\mathcal{G}_\beta) \rangle$ is a dual pair with respect to the bilinear form

$$\langle f, p \rangle = \left\langle \prod_{n \in \mathbb{Z}_+} f_n, \bigoplus_{n \in \mathbb{Z}_+} p_n \right\rangle = \sum_{n \in \mathbb{Z}_+} \langle f_n, p_n \rangle, \quad p \in \Gamma(\mathcal{G}_\beta), \quad f \in \Gamma(\mathcal{G}'_\beta), \quad (1)$$

where $p_n \in \mathcal{G}_\beta^{\widehat{\otimes} n}$ and $f_n \in \mathcal{G}'_\beta{}^{\widehat{\otimes} n} \simeq (\mathcal{G}_\beta^{\widehat{\otimes} n})'$.

By analogy we can construct the dual pairs $\langle \Gamma(E'_\beta), \Gamma(E_\beta) \rangle$ and $\langle \mathcal{P}'(E'_\beta), \mathcal{P}(E'_\beta) \rangle$.

We have the following assertion (see also [10, Proposition 2.1]).

Proposition 1.1. *There exist the linear topological isomorphisms*

$$\Upsilon : \mathcal{P}'(\mathcal{G}'_\beta) \longrightarrow \Gamma(\mathcal{G}'_\beta), \quad \Psi : \mathcal{P}'(E'_\beta) \longrightarrow \Gamma(E'_\beta).$$

Using the Proposition 1.1 and tensor structure of the space $\Gamma(\mathcal{G}'_\beta)$, we extend the map F'^{-1} onto $\Gamma(\mathcal{G}'_\beta)$. First, for elements of total subset of the space $\mathcal{G}'_\beta{}^{\widehat{\otimes} n}$ we define the operator $\mathcal{F}'^{\otimes n} : f^{\otimes n} \mapsto \widehat{f}^{\otimes n}$, $\mathcal{F}'^{\otimes 0} := I_{\mathbb{C}}$, where $\widehat{f}^{\otimes n} := (F'^{-1}f)^{\otimes n}$. Next, we extend the map $\mathcal{F}'^{\otimes n}$ onto whole space $\mathcal{G}'_\beta{}^{\widehat{\otimes} n}$ by linearity and continuity. As a result we obtain the map $\mathcal{F}'^{\otimes n} \in \mathcal{L}(\mathcal{G}'_\beta{}^{\widehat{\otimes} n}, E'_\beta{}^{\widehat{\otimes} n})$. And finally, we define the mapping \mathcal{F}'^{\otimes} by the formula

$$\mathcal{F}'^{\otimes} := (\mathcal{F}'^{\otimes n}) : \Gamma(\mathcal{G}'_\beta) \ni f = (f_n) \quad \mapsto \quad \widehat{f} := (\widehat{f}_n) \in \Gamma(E'_\beta),$$

where $f_n \in \mathcal{G}'_\beta{}^{\widehat{\otimes} n}$, $\widehat{f}_n := \mathcal{F}'^{\otimes n} f_n \in E'_\beta{}^{\widehat{\otimes} n}$.

The following commutative diagram

$$\begin{array}{ccc} \mathcal{P}'(\mathcal{G}'_\beta) & \xrightarrow{\mathcal{F}'^{\otimes}_{\mathcal{P}}} & \mathcal{P}'(E'_\beta) \\ \Upsilon \downarrow & & \uparrow \Psi^{-1} \\ \Gamma(\mathcal{G}'_\beta) & \xrightarrow{\mathcal{F}'^{\otimes}} & \Gamma(E'_\beta) \end{array} \quad (2)$$

uniquely defines the operator $\mathcal{F}'^{\otimes}_{\mathcal{P}} \in \mathcal{L}(\mathcal{P}'(\mathcal{G}'_\beta), \mathcal{P}'(E'_\beta))$.

2 CONVOLUTION OF POLYNOMIAL ULTRADISTRIBUTIONS

Let $g \in \mathcal{G}'_\beta$. Define the shift operator on the space $\mathcal{P}(\mathcal{G}'_\beta)$ with the formula

$$\mathcal{T}_g P(f) := P(f + g), \quad P \in \mathcal{P}(\mathcal{G}'_\beta), \quad f \in \mathcal{G}'_\beta.$$

It is easy to see, that \mathcal{T}_g is a linear continuous operator from the space $\mathcal{P}(\mathcal{G}'_\beta)$ into itself.

Let the symbol \odot_k denotes the (right) k -contraction [6] of symmetric tensor product, i.e., $g^{\otimes k} \odot_k \varphi^{\otimes s} := \langle g, \varphi \rangle^k \varphi^{\otimes (s-k)}$, $k \leq s$, $g \in \mathcal{G}'_\beta$, $\varphi \in \mathcal{G}_\beta$.

Let us show, that for any $g \in \mathcal{G}'_\beta$ the shift operator \mathcal{T}_g acts as follows $P = \sum_n \langle \cdot^{\otimes n}, p_n \rangle \mapsto \mathcal{T}_g P = \sum_n \langle \cdot^{\otimes n}, q_n \rangle$, where $p_n, q_n \in \mathcal{G}_\beta^{\widehat{\otimes} n}$, $n = 0, 1, \dots, m$, $m = \deg P$, and the elements q_n can be obtained by the formula

$$q_n = \sum_{k=0}^{m-n} \frac{(n+k)!}{n!k!} g^{\otimes k} \odot_k p_{n+k}.$$

Without loss of generality we can prove this for polynomials of view $P_{\varphi,m} = \sum_{k=0}^m \langle \cdot^{\otimes k}, \varphi^{\otimes k} \rangle$, where $(1, \varphi, \varphi^{\otimes 2}, \dots, \varphi^{\otimes m}, 0, \dots) \in \Gamma(\mathcal{G}_\beta)$, $\varphi \in \mathcal{G}_\beta$, $m \in \mathbb{Z}_+$.

Indeed,

$$\begin{aligned} \mathcal{T}_g P_{\varphi,m}(f) &= P_{\varphi,m}(f+g) = \sum_{k=0}^m \langle (f+g)^{\otimes k}, \varphi^{\otimes k} \rangle = \sum_{k=0}^m \sum_{n=0}^k C_k^n \langle f^{\otimes n} \widehat{\otimes} g^{\otimes(k-n)}, \varphi^{\otimes k} \rangle \\ &= \sum_{n=0}^m \sum_{k=n}^m C_k^n \langle f^{\otimes n} \widehat{\otimes} g^{\otimes(k-n)}, \varphi^{\otimes k} \rangle = \sum_{n=0}^m \sum_{k=0}^{m-n} C_{n+k}^n \langle f^{\otimes n} \widehat{\otimes} g^{\otimes k}, \varphi^{\otimes(n+k)} \rangle \\ &= \sum_{n=0}^m \sum_{k=0}^{m-n} C_{n+k}^n \langle f^{\otimes n}, \langle g, \varphi \rangle^k \varphi^{\otimes n} \rangle = \sum_{n=0}^m \left\langle f^{\otimes n}, \sum_{k=0}^{m-n} C_{n+k}^n \langle g, \varphi \rangle^k \varphi^{\otimes n} \right\rangle \\ &= \sum_{n=0}^m \left\langle f^{\otimes n}, \sum_{k=0}^{m-n} C_{n+k}^n g^{\otimes k} \odot_k \varphi^{\otimes(n+k)} \right\rangle. \end{aligned}$$

Let us define the convolution of a polynomial ultradistribution $U \in \mathcal{P}'(\mathcal{G}'_\beta)$ and a test function $P \in \mathcal{P}(\mathcal{G}'_\beta)$ with the formula $(U * P)(g) := \langle U, \mathcal{T}_g P \rangle$, $g \in \mathcal{G}'_\beta$, where in the right side there is the pairing of the dual pair $\langle \mathcal{P}'(\mathcal{G}'_\beta), \mathcal{P}(\mathcal{G}'_\beta) \rangle$ (see Proposition 1.1 and formula (1)).

If $U \in \mathcal{P}'(\mathcal{G}'_\beta)$ and $P \in \mathcal{P}(\mathcal{G}'_\beta)$ are represented in the form $U = \times_{n \in \mathbb{Z}_+} \langle u_n, \cdot^{\otimes n} \rangle$ and $P = \sum_{n=0}^m \langle \cdot^{\otimes n}, p_n \rangle$ respectively, then the convolution may be written in the explicit form

$$\begin{aligned} (U * P)(g) &= \sum_{n=0}^m \left\langle u_n, \sum_{k=0}^{m-n} C_{n+k}^n g^{\otimes k} \odot_k p_{n+k} \right\rangle = \sum_{n=0}^m \sum_{k=0}^{m-n} C_{n+k}^n \langle u_n \widehat{\otimes} g^{\otimes k}, p_{n+k} \rangle \\ &= \sum_{k=0}^m \sum_{n=0}^{m-k} C_{n+k}^n \langle g^{\otimes k}, u_n \odot_n p_{n+k} \rangle = \sum_{k=0}^m \left\langle g^{\otimes k}, \sum_{n=0}^{m-k} C_{n+k}^n u_n \odot_n p_{n+k} \right\rangle. \end{aligned} \quad (3)$$

It is clear, that $q_k = \sum_{n=0}^{m-k} C_{n+k}^n u_n \odot_n p_{n+k}$ belongs to the space $\mathcal{G}'_\beta \widehat{\otimes}^k$ for each $k = 0, 1, \dots, m$. It follows, that the convolution $U * P$ is a polynomial from the space $\mathcal{P}(\mathcal{G}'_\beta)$.

For any polynomial ultradistribution $U \in \mathcal{P}'(\mathcal{G}'_\beta)$ the mapping C_U , defined with the formula $C_U : \mathcal{P}(\mathcal{G}'_\beta) \ni P \mapsto U * P \in \mathcal{P}(\mathcal{G}'_\beta)$, is said to be the convolution operator, associated with U .

Let us show, that the composition of two convolution operators C_V and C_U , associated with any $V, U \in \mathcal{P}'(\mathcal{G}'_\beta)$, is a convolution operator, associated with some polynomial ultradistribution, which we denote by $V * U$. Let $V, U \in \mathcal{P}'(\mathcal{G}'_\beta)$ and $P \in \mathcal{P}(\mathcal{G}'_\beta)$ are represented in the form $V = \times_{n \in \mathbb{Z}_+} \langle g^{\otimes n}, \cdot^{\otimes n} \rangle$, $U = \times_{n \in \mathbb{Z}_+} \langle f^{\otimes n}, \cdot^{\otimes n} \rangle$ and $P = \sum_{n=0}^m \langle \cdot^{\otimes n}, \varphi^{\otimes n} \rangle$ respectively, where $f, g \in \mathcal{G}'_\beta$, $\varphi \in \mathcal{G}_\beta$.

Using formula (3), we obtain the following equalities.

$$\begin{aligned} (C_V \circ C_U)(P) &= V * (U * P) = \sum_{n=0}^m \left\langle \cdot^{\otimes n}, \sum_{j=0}^{m-n} C_{n+j}^j g^{\otimes j} \odot_j q_{n+j} \right\rangle \\ &= \sum_{n=0}^m \left\langle \cdot^{\otimes n}, \sum_{j=0}^{m-n} C_{n+j}^j g^{\otimes j} \odot_j \left(\sum_{k=0}^{m-n-j} C_{n+j+k}^k f^{\otimes k} \odot_k \varphi^{\otimes(n+j+k)} \right) \right\rangle \end{aligned}$$

$$\begin{aligned}
&= \sum_{n=0}^m \left\langle \cdot \otimes^n, \sum_{j=0}^{m-n} \sum_{k=0}^{m-n-j} C_{n+j}^j C_{n+j+k}^k \langle g, \varphi \rangle^j \langle f, \varphi \rangle^k \varphi^{\otimes n} \right\rangle \\
&= \sum_{n=0}^m \left\langle \cdot \otimes^n, \sum_{j+k=0}^{m-n} \frac{(n+j+k)!}{n!j!k!} (g^{\otimes j} \hat{\otimes} f^{\otimes k}) \otimes_{j+k} \varphi^{\otimes(n+j+k)} \right\rangle \\
&= \sum_{n=0}^m \left\langle \cdot \otimes^n, \sum_{s=0}^{m-n} \frac{(n+s)!}{n!s!} \sum_{j+k=s} \frac{s!}{j!k!} (g^{\otimes j} \hat{\otimes} f^{\otimes k}) \otimes_s \varphi^{\otimes(n+s)} \right\rangle \\
&= \sum_{n=0}^m \sum_{s=0}^{m-n} \frac{(n+s)!}{n!s!} \sum_{j+k=s} \frac{s!}{j!k!} \left\langle \cdot \otimes^n, (g^{\otimes j} \hat{\otimes} f^{\otimes k}) \otimes_s \varphi^{\otimes(n+s)} \right\rangle \\
&= \sum_{s=0}^m \sum_{n=0}^{m-s} \frac{(n+s)!}{n!s!} \sum_{j+k=s} \frac{s!}{j!k!} \left\langle \cdot \otimes^n \hat{\otimes} g^{\otimes j} \hat{\otimes} f^{\otimes k}, \varphi^{\otimes(n+s)} \right\rangle \\
&= \sum_{s=0}^m \left\langle \sum_{j+k=s} \frac{s!}{j!k!} g^{\otimes j} \hat{\otimes} f^{\otimes k}, \sum_{n=0}^{m-s} C_{n+s}^s (\cdot \otimes^n) \otimes_n \varphi^{\otimes(n+s)} \right\rangle.
\end{aligned}$$

It follows, that the composition $C_V \circ C_U$ is the convolution operator, associated with

$$V * U = \times_{n \in \mathbb{Z}_+} \left\langle \sum_{j+k=n} \frac{n!}{j!k!} g^{\otimes j} \hat{\otimes} f^{\otimes k}, \cdot \otimes^n \right\rangle \in \mathcal{P}'(\mathcal{G}'_\beta). \quad (4)$$

For any polynomial ultradistribution $U \in \mathcal{P}'(\mathcal{G}'_\beta)$ let us define the formal series

$$e^{*U} := \sum_{n \in \mathbb{Z}_+} \frac{1}{n!} U^{*n}, \quad \text{where } U^{*n} := \underbrace{U * \dots * U}_n. \quad (5)$$

Note, that each partial sum of this series belongs to the space $\mathcal{P}'(\mathcal{G}'_\beta)$.

3 HEAT EQUATION ASSOCIATED WITH THE GROSS LAPLACIAN

Let $\{U_t : t \in J\}$ be a family of elements from the space $\mathcal{P}'(\mathcal{G}'_\beta)$, let J be an arbitrary interval $[0, \alpha]$, $\alpha \in \mathbb{R}$, $\alpha \geq 0$. Let us assume, that the function $t \mapsto U_t$ is a continuous map from J into $\mathcal{P}'(\mathcal{G}'_\beta)$. Then the function $t \mapsto \mathcal{F}'_{\mathcal{P}} \otimes U_t$ is continuous map from J into $\mathcal{P}'(E'_\beta)$, where the mapping $\mathcal{F}'_{\mathcal{P}} \otimes$ is defined with formula (2). Therefore, for each $t \in J$ the set $\{\mathcal{F}'_{\mathcal{P}} \otimes U_s : s \in [0, t]\}$ is a compact subset in $\mathcal{P}'(E'_\beta)$. In particular, it is bounded. It follows, that the element

$$\int_0^t \mathcal{F}'_{\mathcal{P}} \otimes U_s ds,$$

belongs to the space $\mathcal{P}'(E'_\beta)$ for each $t \in J$. Hence, in the space $\mathcal{P}'(\mathcal{G}'_\beta)$ there exists a unique element, which we denote $\int_0^t U_s ds$, such that

$$\mathcal{F}'_{\mathcal{P}} \otimes \int_0^t U_s ds = \int_0^t \mathcal{F}'_{\mathcal{P}} \otimes U_s ds.$$

Moreover, the map $E_t = \int_0^t U_s ds$, $t \in J$, is differentiable and satisfies the equality $\frac{\partial}{\partial t} E_t = U_t$.

Let $\{U_t : t \in J\}$ be any described above family of elements from $\mathcal{P}'(\mathcal{G}'_\beta)$. Let us consider the Cauchy problem

$$\begin{cases} \frac{\partial}{\partial t} X_t = U_t * X_t, & t \in J, \\ X_0 = P, & P \in \mathcal{P}(\mathcal{G}'_\beta). \end{cases} \tag{6}$$

Theorem 1. *Cauchy problem (6) has a unique solution in $\mathcal{P}(\mathcal{G}'_\beta)$, which can be presented in the view*

$$X_t = e^{*\int_0^t U_s ds} * P, \quad t \in J, \tag{7}$$

where $e^{*\int_0^t U_s ds}$ is treated in the sense of the formula (5).

Proof. Using Picard’s iteration, the solution X_t of Cauchy problem (6) is written informally in the form (7). Since the polynomial $P \in \mathcal{P}(\mathcal{G}'_\beta)$ has a finite number of addends, a value of $e^{*\int_0^t U_s ds} * P$ depends on some partial sum of the series $e^{*\int_0^t U_s ds}$. Formula (3) implies that solution (7) belongs to the space $\mathcal{P}(\mathcal{G}'_\beta)$. \square

As an application of Theorem 1 we consider the generalized heat equation, associated with the Gross Laplacian.

Let the trace operator τ be defined by

$$\langle \tau, \varphi \hat{\otimes} \psi \rangle := \int_{\mathbb{R}_+^d} \varphi(t) \psi(t) dt, \quad \varphi, \psi \in \mathcal{G}_\beta.$$

It is clear, that $\tau \in \mathcal{L}(\mathcal{G}_\beta^{\hat{\otimes} 2}, \mathbb{C}) = (\mathcal{G}_\beta^{\hat{\otimes} 2})' \simeq \mathcal{G}'_\beta$.

The Gross Laplacian Δ_G by definition (see e.g. [8]) is the following operator

$$\Delta_G : P = \sum_{n=0}^m \langle \cdot^{\otimes n}, \varphi^{\otimes n} \rangle \mapsto \Delta_G P := \sum_{n=0}^{m-2} (n+2)(n+1) \langle \tau, \varphi^{\otimes 2} \rangle \langle \cdot^{\otimes n}, \varphi^{\otimes n} \rangle, \quad \varphi \in \mathcal{G}_\beta.$$

Theorem 2. *The Gross Laplacian Δ_G acts as a convolution operator, i.e.*

$$\frac{1}{2} \Delta_G P = U_\tau * P, \quad P \in \mathcal{P}(\mathcal{G}'_\beta),$$

where U_τ is a polynomial ultradistribution from the space $\mathcal{P}'(\mathcal{G}'_\beta)$, that corresponds to the element $(0, 0, \tau, 0, \dots) \in \Gamma(\mathcal{G}'_\beta)$.

Proof. The polynomial ultradistribution U_τ can be written in the form

$$U_\tau = \times_{n \in \mathbb{Z}_+} \langle u_{\tau, n}, \cdot^{\otimes n} \rangle = (0, 0, \langle \tau, \cdot^{\otimes 2} \rangle, 0, \dots),$$

where $u_{\tau, n} = \tau$ if $n = 2$ and $u_{\tau, n} = 0$ if $n \neq 2$.

Let the polynomial $P \in \mathcal{P}(\mathcal{G}'_\beta)$ be of the form $P = \sum_{n=0}^m \langle \cdot^{\otimes n}, \varphi^{\otimes n} \rangle$, $\varphi \in \mathcal{G}_\beta$. Using equalities (3), we obtain the required result

$$\begin{aligned} U_\tau * P &= \sum_{n=0}^m \left\langle \cdot^{\otimes n}, \sum_{k=0}^{m-n} C_{n+k}^k u_{\tau, k} \otimes_k \varphi^{\otimes(n+k)} \right\rangle = \sum_{n=0}^{m-2} \left\langle \cdot^{\otimes n}, C_{n+2}^2 \tau \otimes_2 \varphi^{\otimes(n+2)} \right\rangle \\ &= \sum_{n=0}^{m-2} C_{n+2}^2 \langle \tau, \varphi^{\otimes 2} \rangle \langle \cdot^{\otimes n}, \varphi^{\otimes n} \rangle = \frac{1}{2} \Delta_G P. \end{aligned}$$

\square

Theorem 3. Cauchy problem

$$\begin{cases} \frac{\partial}{\partial t} X_t = \frac{1}{2} \Delta_G X_t, & t \in J, \\ X_0 = P, & P \in \mathcal{P}(\mathcal{G}'_\beta), \end{cases} \quad (8)$$

for heat equation, associated with the Gross Laplacian, has a unique solution in $\mathcal{P}(\mathcal{G}'_\beta)$ given by

$$X_t = e^{*tU_\tau} * P, \quad t \in J.$$

Proof. Theorem 2 allows us to rewrite the heat equation in the view $\frac{\partial}{\partial t} X_t = U_\tau * X_t$. It follows from Theorem 1 that the Cauchy problem has a unique solution given by

$$X_t = e^{*\int_0^t U_\tau ds} * P = e^{*tU_\tau} * P.$$

We can rewrite it in explicit form. Using formula (4), let us find $(tU_\tau)^{*n}$. For $n = 2$ we obtain

$$(tU_\tau) * (tU_\tau) = \times_{n \in \mathbb{Z}_+} \left\langle t^2 \sum_{j+k=n} \frac{n!}{j!k!} u_{\tau,j} \hat{\otimes} u_{\tau,k}, \cdot^{\otimes n} \right\rangle = (0, 0, 0, 0, \frac{4!}{2!2!} t^2 \langle \tau^{\otimes 2}, \cdot^{\otimes 4} \rangle, 0, \dots),$$

since $u_{\tau,n}$ does not vanish only for $n = 2$. Using the mathematical induction, it is easy to prove that

$$(tU_\tau)^{*n} = (\underbrace{0, \dots, 0}_{2n}, \frac{(2n)!}{2^n} t^n \langle \tau^{\otimes n}, \cdot^{\otimes 2n} \rangle, 0, \dots).$$

It follows

$$\begin{aligned} e^{*tU_\tau} &= \sum_{n \in \mathbb{Z}_+} \frac{1}{n!} (tU_\tau)^{*n} = \sum_{n \in \mathbb{Z}_+} \frac{1}{n!} (\underbrace{0, \dots, 0}_{2n}, \frac{(2n)!}{2^n} t^n \langle \tau^{\otimes n}, \cdot^{\otimes 2n} \rangle, 0, \dots) \\ &= (1, 0, t \langle \tau, \cdot^{\otimes 2} \rangle, 0, 3t^2 \langle \tau^{\otimes 2}, \cdot^{\otimes 4} \rangle, 0, \dots, 0, \underbrace{\frac{(2n)!}{n!} \frac{t^n}{2^n} \langle \tau^{\otimes n}, \cdot^{\otimes 2n} \rangle}_{2n\text{-th place}}, 0, \dots). \end{aligned} \quad (9)$$

It only remains to find the convolution $e^{*tU_\tau} * P$. Let the polynomial $P \in \mathcal{P}(\mathcal{G}'_\beta)$ be written in the form $P = \sum_{n=0}^m \langle \cdot^{\otimes n}, \varphi^{\otimes n} \rangle$, $\varphi \in \mathcal{G}_\beta$. For any $n \in \mathbb{Z}_+$ let us denote $e_{2n} := \frac{(2n)!}{n!} \frac{t^n}{2^n} \tau^{\otimes n}$ and $e_{2n+1} := 0$. Then e^{*tU_τ} can be rewritten as $e^{*tU_\tau} = \times_{n \in \mathbb{Z}_+} \langle e_n, \cdot^{\otimes n} \rangle$. Therefore, we obtain

$$\begin{aligned} e^{*tU_\tau} * P &= \sum_{n=0}^m \left\langle \cdot^{\otimes n}, \sum_{k=0}^{m-n} C_{n+k}^k e_k \otimes_k \varphi^{\otimes(n+k)} \right\rangle = \sum_{n=0}^m \left\langle \cdot^{\otimes n}, \sum_{k=0}^{\lfloor \frac{m-n}{2} \rfloor} C_{n+2k}^{2k} e_{2k} \otimes_{2k} \varphi^{\otimes(n+2k)} \right\rangle \\ &= \sum_{n=0}^m \left\langle \cdot^{\otimes n}, \sum_{k=0}^{\lfloor \frac{m-n}{2} \rfloor} \frac{(n+2k)! (2k)! t^k}{(2k)! n! k! 2^k} \langle \tau^{\otimes k}, \varphi^{\otimes 2k} \rangle \varphi^{\otimes n} \right\rangle \\ &= \sum_{n=0}^m \sum_{k=0}^{\lfloor \frac{m-n}{2} \rfloor} \frac{(n+2k)! t^k}{k! n! 2^k} \langle \tau^{\otimes k}, \varphi^{\otimes 2k} \rangle \langle \cdot^{\otimes n}, \varphi^{\otimes n} \rangle, \end{aligned}$$

where the symbol $\lfloor \cdot \rfloor$ denotes the floor function.

Hence, if the polynomial P from (8) has the form $P = \sum_{n=0}^m \langle \cdot^{\otimes n}, p_n \rangle$, $p_n \in \mathcal{G}'_\beta$, then the solution of Cauchy problem for heat equation associated with the Gross Laplacian can be expressed as

$$X_t = \sum_{n=0}^m \left\langle \cdot^{\otimes n}, \sum_{k=0}^{\lfloor \frac{m-n}{2} \rfloor} \frac{(n+2k)! t^k}{k! n! 2^k} \tau^{\otimes k} \otimes_{2k} p_{n+2k} \right\rangle.$$

□

4 SEMIGROUP GENERATED BY THE GROSS LAPLACIAN

Our next goal is to construct an one-parameter semigroup $\{G_t : t \geq 0\}$ with the infinitesimal generator $\frac{1}{2}\Delta_G$. This semigroup can be formally expressed as $G_t = e^{t\frac{1}{2}\Delta_G}$.

Since $\frac{1}{2}\Delta_G P = U_\tau * P$, results of previous section imply

$$G_t P := \sum_{n=0}^m \sum_{k=0}^{\lfloor \frac{m-n}{2} \rfloor} \frac{(n+2k)!}{k!n!} \frac{t^k}{2^k} \langle \tau^{\otimes k}, \varphi^{\otimes 2k} \rangle \langle \cdot^{\otimes n}, \varphi^{\otimes n} \rangle, \quad (10)$$

where $P = \sum_{n=0}^m \langle \cdot^{\otimes n}, \varphi^{\otimes n} \rangle$, $\varphi \in \mathcal{G}_\beta$.

Proposition 4.1. *The mapping $\mathbb{R}_+ \ni t \mapsto G_t \in \mathcal{L}(\mathcal{P}(\mathcal{G}'_\beta))$, where G_t is defined by formula (10), is a strongly continuous one-parameter semigroup of continuous linear operators from $\mathcal{P}(\mathcal{G}'_\beta)$ into itself with infinitesimal generator $\frac{1}{2}\Delta_G$.*

Proof. Formula (10) can be rewritten as

$$G_t P = P + \sum_{n=0}^{m-2} \sum_{k=1}^{\lfloor \frac{m-n}{2} \rfloor} \frac{(n+2k)!}{k!n!} \frac{t^k}{2^k} \langle \tau^{\otimes k}, \varphi^{\otimes 2k} \rangle \langle \cdot^{\otimes n}, \varphi^{\otimes n} \rangle, \quad (11)$$

therefore the equality $G_0 = I_{\mathcal{P}(\mathcal{G}'_\beta)}$ is clear.

Formulas (4), (9) and the following equalities

$$G_t G_s = e^{t\frac{1}{2}\Delta_G} e^{s\frac{1}{2}\Delta_G} = e^{*tU_\tau} * e^{*sU_\tau} = e^{*(t+s)U_\tau} = e^{(t+s)\frac{1}{2}\Delta_G} = G_{t+s}$$

imply the semigroup property $G_t G_s = G_{t+s}$.

To prove the strong continuity of the semigroup, we need to show that for any $P \in \mathcal{P}(\mathcal{G}'_\beta)$ the function $t \mapsto G_t P$ is continuous. Using representation (11), we obtain

$$\begin{aligned} \limsup_{t \rightarrow 0} \sup_f |G_t P - P| &= \limsup_{t \rightarrow 0} \sup_f \left| \sum_{n=0}^m \sum_{k=1}^{\lfloor \frac{m-n}{2} \rfloor} \frac{(n+2k)!}{k!n!} \frac{t^k}{2^k} \langle \tau^{\otimes k}, \varphi^{\otimes 2k} \rangle \langle f^{\otimes n}, \varphi^{\otimes n} \rangle \right| \\ &\leq \limsup_{t \rightarrow 0} \sup_f \sum_{n=0}^m \sum_{k=1}^{\lfloor \frac{m-n}{2} \rfloor} \frac{(n+2k)!}{k!n!} \frac{|t|^k}{2^k} |\langle \tau^{\otimes k}, \varphi^{\otimes 2k} \rangle| |\langle f^{\otimes n}, \varphi^{\otimes n} \rangle| \\ &= \sum_{n=0}^m \sup_f |\langle f^{\otimes n}, \varphi^{\otimes n} \rangle| \lim_{t \rightarrow 0} \sum_{k=1}^{\lfloor \frac{m-n}{2} \rfloor} \frac{(n+2k)!}{k!n!} \frac{|t|^k}{2^k} |\langle \tau^{\otimes k}, \varphi^{\otimes 2k} \rangle| = 0. \end{aligned}$$

It remains to show that the Gross Laplacian is the generator of the semigroup G_t . Using representation (11), we can write

$$\begin{aligned} \frac{G_t P - P}{t} - \frac{1}{2}\Delta_G P &= \sum_{n=0}^m \sum_{k=1}^{\lfloor \frac{m-n}{2} \rfloor} \frac{(n+2k)!}{k!n!} \frac{t^{k-1}}{2^k} \langle \tau^{\otimes k}, \varphi^{\otimes 2k} \rangle \langle \cdot^{\otimes n}, \varphi^{\otimes n} \rangle \\ &\quad - \sum_{n=0}^{m-2} \frac{(n+2)(n+1)}{2} \langle \tau, \varphi^{\otimes 2} \rangle \langle \cdot^{\otimes n}, \varphi^{\otimes n} \rangle. \end{aligned}$$

Note, that $\lfloor \frac{m-n}{2} \rfloor = 0$ for $n = m - 1$ and for $n = m$. So, we can rewrite the above equality

$$\frac{G_t P - P}{t} - \frac{1}{2} \Delta_G P = \sum_{n=0}^{m-2} \left(\sum_{k=1}^{\lfloor \frac{m-n}{2} \rfloor} \frac{(n+2k)! t^{k-1}}{k!n! 2^k} \langle \tau^{\otimes k}, \varphi^{\otimes 2k} \rangle - \frac{(n+2)(n+1)}{2} \langle \tau, \varphi^{\otimes 2} \rangle \right) \langle \cdot, \otimes^n, \varphi^{\otimes n} \rangle.$$

It is clear that $\frac{(n+2k)! t^{k-1}}{k!n! 2^k} \langle \tau^{\otimes k}, \varphi^{\otimes 2k} \rangle = \frac{(n+2)(n+1)}{2} \langle \tau, \varphi^{\otimes 2} \rangle$ with $k = 1$, therefore

$$\frac{G_t P - P}{t} - \frac{1}{2} \Delta_G P = \sum_{n=0}^{m-2} \left(\sum_{k=2}^{\lfloor \frac{m-n}{2} \rfloor} \frac{(n+2k)! t^{k-1}}{k!n! 2^k} \langle \tau^{\otimes k}, \varphi^{\otimes 2k} \rangle \right) \langle \cdot, \otimes^n, \varphi^{\otimes n} \rangle.$$

Note, that $\lfloor \frac{m-n}{2} \rfloor = 1$ for $n = m - 2$ and for $n = m - 3$. So, we obtain

$$\frac{G_t P - P}{t} - \frac{1}{2} \Delta_G P = \sum_{n=0}^{m-4} \left(\sum_{k=2}^{\lfloor \frac{m-n}{2} \rfloor} \frac{(n+2k)! t^{k-1}}{k!n! 2^k} \langle \tau^{\otimes k}, \varphi^{\otimes 2k} \rangle \right) \langle \cdot, \otimes^n, \varphi^{\otimes n} \rangle.$$

From the above formula we can derive the required result

$$\begin{aligned} \limsup_{t \rightarrow 0} \sup_f \left| \frac{G_t P(f) - P(f)}{t} - \frac{1}{2} \Delta_G P(f) \right| \\ \leq \sum_{n=0}^{m-4} \sup_f |\langle f^{\otimes n}, \varphi^{\otimes n} \rangle| \lim_{t \rightarrow 0} \sum_{k=2}^{\lfloor \frac{m-n}{2} \rfloor} \frac{(n+2k)! |t|^{k-1}}{k!n! 2^k} |\langle \tau^{\otimes k}, \varphi^{\otimes 2k} \rangle| = 0. \end{aligned}$$

□

Corollary 4.1. *Cauchy problem*

$$\begin{cases} \frac{\partial}{\partial t} X_t = \frac{1}{2} \Delta_G X_t, & t \in J, \\ X_0 = P, & P \in \mathcal{P}(\mathcal{G}'_\beta), \end{cases}$$

for heat equation associated with the Gross Laplacian has a unique solution in $\mathcal{P}(\mathcal{G}'_\beta)$ given by

$$X_t = G_t P, \quad t \in J.$$

REFERENCES

- [1] Berezanskii Yu.M., Samoilenko Yu.S. *Nuclear spaces of functions of infinitely many variables*. Ukrainian Math. J. 1973, **25** (6), 599–609. doi:10.1007/BF01090792 (translation of Ukrain. Mat. Zh. 1973, **25** (6), 723–737. (in Russian))
- [2] Chung D.M., Ji U.C. *Some Cauchy problems in white noise analysis and associated semigroups of operators*. Stochastic Anal. Appl. 1999, **17**, (1), 1–22. doi:10.1080/07362999908809585
- [3] Dineen S. *Complex analysis on infinite dimensional spaces*. Springer, Berlin-Göttingen-Heidelberg, 1999.
- [4] Gross L. *Potential theory on Hilbert space*. J. Funct. Anal. 1967, **1** (2), 123–181. doi:10.1016/0022-1236(67)90030-4
- [5] Hida T. *Analysis of Brownian Functionals*. Carleton Math. Lect. Notes, Vol. 13. Carleton Univ. Press, Ottawa, 1975.

- [6] Ji U.C., Obata N. Admissible white noise operators and their quantum white noise derivatives. In: Heyer H. et al. (Ed.) Infinite dimensional harmonic analysis III. World Sci. Publ., RiverEdge, NJ, 2005. P. 213–232.
- [7] Komatsu H. An introduction to the theory of generalized functions University Publ., Tokyo, 2000.
- [8] Kuo H. White Noise Distribution Theory. CRC Press, Boca Ration, FL, 1996.
- [9] Lee J.-S.K. *Heat Equation In White Noise Analysis*. J. Korean Math. Soc. 1996, **33** (3), 541–555.
- [10] Lopushansky O.V., Sharyn S.V. *Polynomial ultradistributions on cone \mathbb{R}_+^d* . Topology 2009, **48** (2-4), 80–90. doi:10.1016/j.top.2009.11.005
- [11] Schaefer H.H. Topological Vector Spaces. Springer Verlag, New York, 1999.
- [12] Sharyn S. *Joint functional calculus in algebra of polynomial tempered distributions*. Methods Funct. Anal. Topology 2016, **22** (1), 62–73.
- [13] Sharyn S.V. *Paley–Wiener-type theorem for polynomial ultradifferentiable functions*. Carpathian Math. Publ. 2015, **7** (2), 271–279. doi:10.15330/cmp.7.2.271-279

Received 20.11.2016

Шарин С.В. Застосування функціонального числення до розв'язання задачі Коші для нескінченновимірної рівняння теплопровідності // Карпатські матем. публ. — 2016. — Т.8, №2. — С. 313–322.

У цій роботі ми вивчаємо нескінченновимірне рівняння теплопровідності, породжене лапласіаном Гросса. Використовуючи метод функціонального числення, ми отримуємо розв'язок відповідної задачі Коші у просторі поліноміальних ультрадиференційовних функцій. Також розглянуто напівгруповий підхід розв'язання такої задачі.

Ключові слова і фрази: нескінченновимірне рівняння теплопровідності, лапласіан Гросса, простір поліноміальних ультрадиференційовних функцій, простір поліноміальних ультра-розподілів.